

**TABLE 10.2** Prevalence and Its Standard Error for Eating Disorder by Gender

	MALE		FEMALE		TOTAL	
	%	(SE)	%	(SE)	%	(SE)
<i>I. Lifetime prevalence</i>						
Anorexia nervosa	.3*	(.1)	.9*	(.3)	.6	(.2)
Bulimia nervosa	.5*	(.3)	1.5*	(.3)	1.0	(.2)
Binge eating disorder	2.0*	(.5)	3.5*	(.5)	2.8	(.4)
Subthreshold binge eating disorder	1.9*	(.5)	.6*	(.1)	1.2	(.2)
Any binge eating	4.0	(.7)	4.9	(.6)	4.5	(.4)
<i>II. Twelve-month prevalence<sup>†</sup></i>						
Bulimia nervosa	.1*	(.1)	.5*	(.2)	.3	(.1)
Binge eating disorder	.8*	(.3)	1.6*	(.2)	1.2	(.2)
Subthreshold binge eating disorder	.8	(.3)	.4	(.1)	.6	(.2)
Any binge eating	1.7	(.4)	2.5	(.3)	2.1	(.2)
(n)	(1,220)		(1,760)		(2,980)	

Abbreviations: SE, standard error.

\*Significant sex difference based on a .05 level, 2-sided test.

<sup>†</sup>None of the respondents met criteria for 12-month anorexia nervosa.

Source: Hudson et al. (2007, p. 350), with permission from Elsevier.