

**TABLE 8.4**   Lifetime Prevalence in Terms of Percentage and Standard Error (SE)

|  | ANY ANXIETY DISORDER |       |
|--|----------------------|-------|
|  | %                    | (SE)  |
| <i>I. WHO Region: Pan American Health</i>      |                      |       |
| Colombia                                       | 25.3                 | (1.4) |
| Mexico   | 14.3                 | (0.9) |
| United States                                  | 31.0                 | (1.0) |
| <i>II. WHO Region: African Regional Office</i> |                      |       |
| Nigeria  | 6.5                  | (0.9) |
| South Africa                                   | 15.8                 | (0.8) |
| <i>III. WHO Region: Eastern Mediterranean</i>  |                      |       |
| Lebanon  | 16.7                 | (1.6) |
| <i>IV. WHO Region: European Regional</i>       |                      |       |
| Belgium  | 13.1                 | (1.9) |
| France   | 22.3                 | (1.4) |
| Germany  | 14.6                 | (1.5) |
| Israel   | 5.2                  | (0.3) |
| Italy  | 11.0                 | (0.9) |
| Netherlands                                    | 15.9                 | (1.1) |
| Spain  | 9.9                  | (1.1) |
| Ukraine  | 10.9                 | (0.8) |
| <i>V. WHO Region: Western Pacific Region</i>   |                      |       |
| PRC  | 4.8                  | (0.7) |
| Japan  | 6.9                  | (0.6) |
| New Zealand                                    | 24.6                 | (0.7) |

Source: Kessler et al. (2009).