

TABLE 9.1 A Portion of the *t* Table Adapted From Table B.2 in Appendix B

df	<i>Proportion in One Tail</i>					
	<i>.25</i>	<i>.10</i>	<i>.05</i>	<i>.025</i>	<i>.01</i>	<i>.005</i>
	<i>Proportion in Two Tails Combined</i>					
	<i>.50</i>	<i>.20</i>	<i>.10</i>	<i>.05</i>	<i>.02</i>	<i>.01</i>
1	1.000	3.078	6.314	12.706	31.821	63.657
2	0.816	1.886	2.920	4.303	6.965	9.925
3	0.765	1.638	2.353	3.182	4.541	5.841
4	0.741	1.533	2.132	2.776	3.747	4.604
5	0.727	1.476	2.015	2.571	3.365	4.032
6	0.718	1.440	1.943	2.447	3.143	3.707
7	0.711	1.415	1.895	2.365	2.998	3.499
8	0.706	1.397	1.860	2.306	2.896	3.355
9	0.703	1.383	1.833	2.282	2.821	3.250
10	0.700	1.372	1.812	2.228	2.764	3.169

Source: Table III of R. A. Fisher and F. Yates, *Statistical Tables for Biological, Agricultural and Medical Research*, 6th ed. London: Longman Group Ltd., 1974 (previously published by Oliver and Boyd Ltd., Edinburgh). Adapted and reprinted with permission of Addison Wesley Longman.